

Metastability

in a class of parabolic SPDEs

Nils Berglund

MAPMO, Université d'Orléans

CNRS, UMR 6628 et Fédération Denis Poisson

www.univ-orleans.fr/mapmo/membres/berglund

Collaborators:

[Florent Barret](#), Ecole Polytechnique, Palaiseau

[Bastien Fernandez](#), CPT, CNRS, Marseille

[Barbara Gentz](#), University of Bielefeld

Grenoble, 2 décembre 2010

Metastability in physics

Examples:

- Supercooled liquid
- Supersaturated gas
- Wrongly magnetised ferromagnet




Metastability in physics

Examples:

- Supercooled liquid
- Supersaturated gas
- Wrongly magnetised ferromagnet



- ▷ Near first-order phase transition
- ▷ Nucleation implies crossing energy barrier



Metastability in stochastic lattice models


- ▷ Lattice: $\Lambda \subset \mathbb{Z}^d$
- ▷ Configuration space: $\mathcal{X} = S^\Lambda$, S finite set (e.g. $\{-1, 1\}$)
- ▷ Hamiltonian: $H : \mathcal{X} \rightarrow \mathbb{R}$ (e.g. Ising or lattice gas)
- ▷ Gibbs measure: $\mu_\beta(x) = e^{-\beta H(x)} / Z_\beta$
- ▷ Dynamics: Markov chain with invariant measure μ_β
(e.g. Metropolis: Glauber or Kawasaki)

Metastability in stochastic lattice models

- ▷ Lattice: $\Lambda \subset \mathbb{Z}^d$
- ▷ Configuration space: $\mathcal{X} = S^\Lambda$, S finite set (e.g. $\{-1, 1\}$)
- ▷ Hamiltonian: $H : \mathcal{X} \rightarrow \mathbb{R}$ (e.g. Ising or lattice gas)
- ▷ Gibbs measure: $\mu_\beta(x) = e^{-\beta H(x)} / Z_\beta$
- ▷ Dynamics: Markov chain with invariant measure μ_β (e.g. Metropolis: Glauber or Kawasaki)

Results (for $\beta \gg 1$) on

- Transition time between $+$ and $-$ or empty and full configuration
- Transition path
- Shape of critical droplet



- ▷ Frank den Hollander, *Metastability under stochastic dynamics*, Stochastic Process. Appl. **114** (2004), 1–26.
- ▷ Enzo Olivieri and Maria Eulália Vares, *Large deviations and metastability*, Cambridge University Press, Cambridge, 2005.

Reversible diffusion

$$dx_t = -\nabla V(x_t) dt + \sqrt{2\varepsilon} dW_t$$

- ▷ $V : \mathbb{R}^d \rightarrow \mathbb{R}$: potential, growing at infinity
- ▷ W_t : d -dim Brownian motion on $(\Omega, \mathcal{F}, \mathbb{P})$

Reversible w.r.t.

invariant measure:

$$\mu_\varepsilon(dx) = \frac{e^{-V(x)/\varepsilon}}{Z_\varepsilon} dx$$

(detailed balance)

Reversible diffusion


$$dx_t = -\nabla V(x_t) dt + \sqrt{2\varepsilon} dW_t$$

- ▷ $V : \mathbb{R}^d \rightarrow \mathbb{R}$: potential, growing at infinity
- ▷ W_t : d -dim Brownian motion on $(\Omega, \mathcal{F}, \mathbb{P})$

Reversible w.r.t.
invariant measure:

$$\mu_\varepsilon(dx) = \frac{e^{-V(x)/\varepsilon}}{Z_\varepsilon} dx$$

(detailed balance)



τ_y^x : first-hitting time of small ball $\mathcal{B}_\varepsilon(y)$, starting in x
“Eyring–Kramers law” (Eyring 1935, Kramers 1940)

- Dim 1: $\mathbb{E}[\tau_y^x] \simeq \frac{2\pi}{\sqrt{V''(x)|V''(z)|}} e^{[V(z)-V(x)]/\varepsilon}$
- Dim ≥ 2 : $\mathbb{E}[\tau_y^x] \simeq \frac{2\pi}{|\lambda_1(z)|} \sqrt{\frac{|\det(\nabla^2 V(z))|}{\det(\nabla^2 V(x))}} e^{[V(z)-V(x)]/\varepsilon}$

Towards a proof of Kramers' law

- Large deviations (Wentzell & Freidlin 1969):

$$\lim_{\varepsilon \rightarrow 0} \varepsilon \log(\mathbb{E}[\tau_y^x]) = V(z) - V(x)$$

- Analytic (Helffer, Sjöstrand 85, Miclo 95, Mathieu 95, Kolokoltsov 96, . . .):
low-lying spectrum of generator
- Potential theory/variational (Bovier, Eckhoff, Gaynard, Klein 2004):

$$\mathbb{E}[\tau_y^x] = \frac{2\pi}{|\lambda_1(z)|} \sqrt{\frac{|\det(\nabla^2 V(z))|}{\det(\nabla^2 V(x))}} e^{[V(z)-V(x)]/\varepsilon} \left[1 + \mathcal{O}(\varepsilon^{1/2} |\log \varepsilon|^{1/2}) \right]$$

and similar asymptotics for eigenvalues of generator

- Witten complex (Helffer, Klein, Nier 2004):
full asymptotic expansion of prefactor
- Distribution of τ_y^x (Day 1983, Bovier *et al* 2005):

$$\lim_{\varepsilon \rightarrow 0} \mathbb{P}\left\{ \tau_y^x > t \mathbb{E}[\tau_y^x] \right\} = e^{-t}$$

Allen-Cahn equation

$$\partial_t u(x, t) = \partial_{xx} u(x, t) + f(u(x, t)) + \text{noise}$$

where e.g. $f(u) = u - u^3$

$x \in [0, L]$, $u(x, t) \in \mathbb{R}$ represents e.g. magnetisation

- Periodic b.c.
- Neumann b.c. $\partial_x u(0, t) = \partial_x u(L, t) = 0$

Noise: weak, white in space and time

Allen-Cahn equation

$$\partial_t u(x, t) = \partial_{xx} u(x, t) + f(u(x, t)) + \text{noise}$$

where e.g. $f(u) = u - u^3$

$x \in [0, L]$, $u(x, t) \in \mathbb{R}$ represents e.g. magnetisation

- Periodic b.c.
- Neumann b.c. $\partial_x u(0, t) = \partial_x u(L, t) = 0$

Noise: weak, white in space and time

Deterministic system is gradient

$$\partial_{xx} u + u - u^3 = -\frac{\delta V}{\delta u}$$

$$V[u] = \int_0^L \left[\frac{1}{2} u'(x)^2 - \frac{1}{2} u(x)^2 + \frac{1}{4} u(x)^4 \right] dx \quad \rightarrow \quad \min$$

Stationary solutions

$$u''(x) = -u(x) + u(x)^3 = -\frac{d}{dx} \left[\text{wavy line} \right]$$

Stationary solutions

$$u''(x) = -u(x) + u(x)^3 = -\frac{d}{dx} \left[\text{graph of } V(u) \right]$$

- $u_{\pm}(x) \equiv \pm 1$: global minima of V , stable
- $u_0(x) \equiv 0$: unstable
- **Neumann b.c**: for $k = 1, 2, \dots$, if $L > \pi k$,


$$u_{k,\pm}(x) = \pm \sqrt{\frac{2m}{m+1}} \operatorname{sn}\left(\frac{kx}{\sqrt{m+1}} + K(m), m\right) \quad 2k\sqrt{m+1}K(m) = L$$

Stationary solutions

$$u''(x) = -u(x) + u(x)^3 = -\frac{d}{dx} \left[\text{graph of potential } V(u) \right]$$

- $u_{\pm}(x) \equiv \pm 1$: global minima of V , stable
- $u_0(x) \equiv 0$: unstable
- **Neumann b.c**: for $k = 1, 2, \dots$, if $L > \pi k$,
 $u_{k,\pm}(x) = \pm \sqrt{\frac{2m}{m+1}} \operatorname{sn}\left(\frac{kx}{\sqrt{m+1}} + K(m), m\right)$

$$2k\sqrt{m+1}K(m) = L$$



Stability of stationary solutions


Linearisation at $u(x)$: $\partial_t \varphi = A[u] \varphi$, $A[u] = \frac{d^2}{dx^2} + 1 - 3u(x)^2$

Stability of stationary solutions

Linearisation at $u(x)$: $\partial_t \varphi = A[u]\varphi$, $A[u] = \frac{d^2}{dx^2} + 1 - 3u(x)^2$

- $u_{\pm}(x) \equiv \pm 1$: eigenvalues $-(2 + (\pi k/L)^2)$, $k \in \mathbb{N}$
- $u_0(x) \equiv 0$: eigenvalues $1 - (\pi k/L)^2$, $k \in \mathbb{N}$

Number of positive eigenvalues:



Allen-Cahn equation with noise

$$\dot{u}_t(x) = \Delta u_t(x) + f(u_t(x)) + \sqrt{2\varepsilon} \ddot{W}_{tx} \quad (\Delta \equiv \partial_{xx}, f(u) = u - u^3)$$

\ddot{W}_{tx} : space-time white noise (formal derivative of Brownian sheet)

Allen-Cahn equation with noise

$$\dot{u}_t(x) = \Delta u_t(x) + f(u_t(x)) + \sqrt{2\varepsilon} \ddot{W}_{tx} \quad (\Delta \equiv \partial_{xx}, f(u) = u - u^3)$$

\ddot{W}_{tx} : space-time white noise (formal derivative of Brownian sheet)

Construction of mild solution:

$$1. \dot{u}_t = \Delta u_t \quad \Rightarrow \quad u_t = e^{\Delta t} u_0$$

$$\text{where } e^{\Delta t} \cos(k\pi x/L) = e^{-(k\pi/L)^2 t} \cos(k\pi x/L)$$

Allen-Cahn equation with noise

$$\dot{u}_t(x) = \Delta u_t(x) + f(u_t(x)) + \sqrt{2\varepsilon} \ddot{W}_{tx} \quad (\Delta \equiv \partial_{xx}, f(u) = u - u^3)$$

\ddot{W}_{tx} : space-time white noise (formal derivative of Brownian sheet)

Construction of mild solution:

1. $\dot{u}_t = \Delta u_t \quad \Rightarrow \quad u_t = e^{\Delta t} u_0$
where $e^{\Delta t} \cos(k\pi x/L) = e^{-(k\pi/L)^2 t} \cos(k\pi x/L)$

2. $\dot{u}_t = \Delta u_t + \sqrt{2\varepsilon} \ddot{W}_{tx}$
 $\Rightarrow \quad u_t = e^{\Delta t} u_0 + \underbrace{\sqrt{2\varepsilon} \int_0^t e^{\Delta(t-s)} \dot{W}_x(ds)}_{\doteq w_t(x)}$

$$w_t(x) = \sum_k \int_0^t e^{-(k\pi/L)^2(t-s)} dW_s^{(k)} \cos(k\pi x/L)$$

Allen-Cahn equation with noise

$$\dot{u}_t(x) = \Delta u_t(x) + f(u_t(x)) + \sqrt{2\varepsilon} \ddot{W}_{tx} \quad (\Delta \equiv \partial_{xx}, f(u) = u - u^3)$$

\ddot{W}_{tx} : space-time white noise (formal derivative of Brownian sheet)

Construction of mild solution:

$$1. \dot{u}_t = \Delta u_t \quad \Rightarrow \quad u_t = e^{\Delta t} u_0$$

$$\text{where } e^{\Delta t} \cos(k\pi x/L) = e^{-(k\pi/L)^2 t} \cos(k\pi x/L)$$

$$2. \dot{u}_t = \Delta u_t + \sqrt{2\varepsilon} \ddot{W}_{tx}$$

$$\Rightarrow \quad u_t = e^{\Delta t} u_0 + \underbrace{\sqrt{2\varepsilon} \int_0^t e^{\Delta(t-s)} \dot{W}_x(ds)}_{\doteq w_t(x)}$$

$$w_t(x) = \sum_k \int_0^t e^{-(k\pi/L)^2(t-s)} dW_s^{(k)} \cos(k\pi x/L)$$

$$3. \dot{u}_t = \Delta u_t + \sqrt{2\varepsilon} \ddot{W}_{tx} + f(u_t)$$

$$\Rightarrow \quad u_t = e^{\Delta t} u_0 + \sqrt{2\varepsilon} w_t + \int_0^t e^{\Delta(t-s)} f(u_s) ds =: F_t[u]$$

\Rightarrow Existence and a.s. uniqueness (Faris & Jona-Lasinio 1982)

Allen-Cahn equation with noise

$$\dot{u}_t(x) = \Delta u_t(x) + f(u_t(x)) + \sqrt{2\varepsilon} \ddot{W}_{tx} \quad (\Delta \equiv \partial_{xx}, f(u) = u - u^3)$$

Fourier variables: $u_t(x) = \sum_{k=-\infty}^{\infty} z_k(t) e^{i\pi kx/L}$

$$\Rightarrow dz_k = -\lambda_k z_k dt - \sum_{k_1+k_2+k_3=k} z_{k_1} z_{k_2} z_{k_3} dt + \sqrt{2\varepsilon} dW_t^{(k)}$$

where $\lambda_k = -1 + (\pi k/L)^2$

Allen-Cahn equation with noise

$$\dot{u}_t(x) = \Delta u_t(x) + f(u_t(x)) + \sqrt{2\varepsilon} \dot{W}_{tx} \quad (\Delta \equiv \partial_{xx}, f(u) = u - u^3)$$

Fourier variables: $u_t(x) = \sum_{k=-\infty}^{\infty} z_k(t) e^{i\pi kx/L}$

$$\Rightarrow dz_k = -\lambda_k z_k dt - \sum_{k_1+k_2+k_3=k} z_{k_1} z_{k_2} z_{k_3} dt + \sqrt{2\varepsilon} dW_t^{(k)}$$

where $\lambda_k = -1 + (\pi k/L)^2$

Energy functional:

$$\frac{1}{L} V[u] = \frac{1}{2} \sum_{k=-\infty}^{\infty} \lambda_k |z_k|^2 + \frac{1}{4} \sum_{k_1+k_2+k_3+k_4=0} z_{k_1} z_{k_2} z_{k_3} z_{k_4}$$

The question

How long does the system take to get from $u_-(x) \equiv -1$ to (a neighbourhood of) $u_+(x) \equiv 1$?

Metastability: Time of order $e^{\text{const}/\varepsilon}$
(rate of order $e^{-\text{const}/\varepsilon}$)

The question

How long does the system take to get from $u_-(x) \equiv -1$ to (a neighbourhood of) $u_+(x) \equiv 1$?

Metastability: Time of order $e^{\text{const}/\varepsilon}$
(rate of order $e^{-\text{const}/\varepsilon}$)

We seek constants ΔW (activation energy), Γ_0 and α such that the random transition time τ satisfies

$$\mathbb{E}[\tau] = \Gamma_0^{-1} e^{\Delta W/\varepsilon} [1 + \mathcal{O}(\varepsilon^\alpha)]$$

The question

How long does the system take to get from $u_-(x) \equiv -1$ to (a neighbourhood of) $u_+(x) \equiv 1$?

Metastability: Time of order $e^{\text{const}/\varepsilon}$
(rate of order $e^{-\text{const}/\varepsilon}$)

We seek constants ΔW (activation energy), Γ_0 and α such that the random transition time τ satisfies

$$\mathbb{E}[\tau] = \Gamma_0^{-1} e^{\Delta W/\varepsilon} [1 + \mathcal{O}(\varepsilon^\alpha)]$$

Large deviations (**Faris & Jona-Lasinio 1982**)

$$\triangleright L \leq \pi : \Delta W = V[u_0] - V[u_-] = \frac{L}{4}$$

$$\triangleright L > \pi : \Delta W = V[u_{1,\pm}] - V[u_-] = \frac{1}{3\sqrt{1+m}} \left[8 E(m) - \frac{(1-m)(3m+5)}{1+m} K(m) \right]$$

Formal computation for Allen-Cahn (R.S. Maier, D. Stein, 01)

Case $L < \pi$:

$$\text{Kramers: } \Gamma_0 \simeq \frac{|\lambda_1(u_0)|}{2\pi} \sqrt{\frac{\det(\nabla^2 V[u_-])}{|\det(\nabla^2 V[u_0])|}}$$

Eigenvalues at $V[u_-] \equiv -1$: $\mu_k = 2 + (\pi k/L)^2$

Eigenvalues at $V[u_0] \equiv 0$: $\lambda_k = -1 + (\pi k/L)^2$

Formal computation for Allen-Cahn (R.S. Maier, D. Stein, 01)

Case $L < \pi$:

$$\text{Kramers: } \Gamma_0 \simeq \frac{|\lambda_1(u_0)|}{2\pi} \sqrt{\frac{\det(\nabla^2 V[u_-])}{|\det(\nabla^2 V[u_0])|}}$$

Eigenvalues at $V[u_-] \equiv -1$: $\mu_k = 2 + (\pi k/L)^2$

Eigenvalues at $V[u_0] \equiv 0$: $\lambda_k = -1 + (\pi k/L)^2$

$$\text{Thus formally } \Gamma_0 \simeq \frac{|\lambda_0|}{2\pi} \sqrt{\prod_{k=0}^{\infty} \frac{\mu_k}{|\lambda_k|}} = \frac{1}{2^{3/4}\pi} \sqrt{\frac{\sinh(\sqrt{2}L)}{\sin L}}$$

Case $L > \pi$: Spectral determinant computed by Gelfand's method

Formal computation for Allen-Cahn (R.S. Maier, D. Stein, 01)

Case $L < \pi$:

$$\text{Kramers: } \Gamma_0 \simeq \frac{|\lambda_1(u_0)|}{2\pi} \sqrt{\frac{\det(\nabla^2 V[u_-])}{|\det(\nabla^2 V[u_0])|}}$$

Eigenvalues at $V[u_-] \equiv -1$: $\mu_k = 2 + (\pi k/L)^2$

Eigenvalues at $V[u_0] \equiv 0$: $\lambda_k = -1 + (\pi k/L)^2$

$$\text{Thus formally } \Gamma_0 \simeq \frac{|\lambda_0|}{2\pi} \sqrt{\prod_{k=0}^{\infty} \frac{\mu_k}{|\lambda_k|}} = \frac{1}{2^{3/4}\pi} \sqrt{\frac{\sinh(\sqrt{2}L)}{\sin L}}$$

Case $L > \pi$: Spectral determinant computed by Gelfand's method

Problems:

1. What happens when $L \rightarrow \pi_-$? (bifurcation)
2. Is the formal computation correct in infinite dimension?

Theorem: [Barret, B., Gentz 2010]

$$\mathbb{E}[\tau] = \Gamma_0^{-1} e^{\Delta W/\varepsilon} \left[1 + \mathcal{O}(\varepsilon^{1/4} |\log \varepsilon|^{5/4}) \right]$$

Theorem: [Barret, B., Gentz 2010]

$$\mathbb{E}[\tau] = \Gamma_0^{-1} e^{\Delta W/\varepsilon} \left[1 + \mathcal{O}(\varepsilon^{1/4} |\log \varepsilon|^{5/4}) \right]$$

$L < \pi$:

$$\Gamma_0 = \frac{1}{2^{3/4} \pi} \sqrt{\frac{\sinh(\sqrt{2}L)}{\sin L}} \sqrt{\frac{\lambda_1}{\lambda_1 + \sqrt{3\varepsilon/4L}}} \Psi_+ \left(\frac{\lambda_1}{\sqrt{3\varepsilon/4L}} \right)$$

where $\lambda_1 = -1 + (\pi/L)^2$ and

$$\Psi_+(\alpha) = \sqrt{\frac{\alpha(1+\alpha)}{8\pi}} e^{\alpha^2/16} K_{1/4} \left(\frac{\alpha^2}{16} \right)$$

Theorem: [Barret, B., Gentz 2010]

$$\mathbb{E}[\tau] = \Gamma_0^{-1} e^{\Delta W/\varepsilon} \left[1 + \mathcal{O}(\varepsilon^{1/4} |\log \varepsilon|^{5/4}) \right]$$

$L < \pi$:

$$\Gamma_0 = \frac{1}{2^{3/4} \pi} \sqrt{\frac{\sinh(\sqrt{2}L)}{\sin L}} \sqrt{\frac{\lambda_1}{\lambda_1 + \sqrt{3\varepsilon/4L}}} \Psi_+ \left(\frac{\lambda_1}{\sqrt{3\varepsilon/4L}} \right)$$

where $\lambda_1 = -1 + (\pi/L)^2$ and

$$\Psi_+(\alpha) = \sqrt{\frac{\alpha(1+\alpha)}{8\pi}} e^{\alpha^2/16} K_{1/4} \left(\frac{\alpha^2}{16} \right)$$

In particular,

$$\lim_{L \rightarrow \pi^-} \Gamma_0 = \frac{\Gamma(1/4)}{2(3\pi^7)^{1/4}} \sqrt{\sinh(\sqrt{2}\pi)} \varepsilon^{-1/4}$$

Theorem: [Barret, B., Gentz 2010]

$$\mathbb{E}[\tau] = \Gamma_0^{-1} e^{\Delta W/\varepsilon} \left[1 + \mathcal{O}(\varepsilon^{1/4} |\log \varepsilon|^{5/4}) \right]$$

$L < \pi$:

$$\Gamma_0 = \frac{1}{2^{3/4} \pi} \sqrt{\frac{\sinh(\sqrt{2}L)}{\sin L}} \sqrt{\frac{\lambda_1}{\lambda_1 + \sqrt{3\varepsilon/4L}}} \Psi_+ \left(\frac{\lambda_1}{\sqrt{3\varepsilon/4L}} \right)$$

where $\lambda_1 = -1 + (\pi/L)^2$ and


$$\Psi_+(\alpha) = \sqrt{\frac{\alpha(1+\alpha)}{8\pi}} e^{\alpha^2/16} K_{1/4} \left(\frac{\alpha^2}{16} \right)$$

In particular,

$$\lim_{L \rightarrow \pi^-} \Gamma_0 = \frac{\Gamma(1/4)}{2(3\pi^7)^{1/4}} \sqrt{\sinh(\sqrt{2}\pi)} \varepsilon^{-1/4}$$

Similar expression for $L > \pi$

(product of eigenvalues computed using path-integral techniques, cf. Maier and Stein)



Spectral Galerkin approximation

$$u_t^d(x) = \sum_{k=-d}^d z_k(t) e^{i\pi kx/L}$$

$z_k(t)$ solution of finite-dimensional SDE

$$dz_k(t) = -\nabla V(z(t)) dt + \sqrt{2\varepsilon} dW_k(t)$$

Spectral Galerkin approximation

$$u_t^d(x) = \sum_{k=-d}^d z_k(t) e^{i\pi kx/L}$$

$z_k(t)$ solution of finite-dimensional SDE

$$dz_k(t) = -\nabla V(z(t)) dt + \sqrt{2\varepsilon} dW_k(t)$$

Theorem: [D. Blömker, A. Jentzen 2009]

Assume

$$\sup_{d \in \mathbb{N}} \sup_{0 \leq t \leq T} \|u_t^d(\omega)\|_{L^\infty} < \infty \quad \forall \omega \in \Omega$$

Then

$$\sup_{0 \leq t \leq T} \|u_t(\omega) - u_t^d(\omega)\|_{L^\infty} < Z(\omega) d^{-\gamma} \quad \forall \omega \in \Omega$$

for all $0 < \gamma < 1/2$, with Z a.s. finite r.v.

Spectral Galerkin approximation

$$u_t^d(x) = \sum_{k=-d}^d z_k(t) e^{i\pi kx/L}$$

$z_k(t)$ solution of finite-dimensional SDE

$$dz_k(t) = -\nabla V(z(t)) dt + \sqrt{2\varepsilon} dW_k(t)$$

Theorem: [D. Blömker, A. Jentzen 2009]

Assume

$$\sup_{d \in \mathbb{N}} \sup_{0 \leq t \leq T} \|u_t^d(\omega)\|_{L^\infty} < \infty \quad \forall \omega \in \Omega$$

Then

$$\sup_{0 \leq t \leq T} \|u_t(\omega) - u_t^d(\omega)\|_{L^\infty} < Z(\omega) d^{-\gamma} \quad \forall \omega \in \Omega$$

for all $0 < \gamma < 1/2$, with Z a.s. finite r.v.

Rem: [Liu, 2003] similar result for $\|u\|^2 = \sum_k (1 + k^2)^r |z_k|^2$, $r < \frac{1}{2}$

Potential theory

$$\text{SDE: } dz_t = -\nabla V(z_t) dt + \sqrt{2\varepsilon} dW_t$$

$$\text{Equilibrium potential : } h_{A,B}(x) = \mathbb{P}^x\{\tau_A < \tau_B\}$$

Then

$$\mathbb{E}[\tau_B^x] \simeq \frac{\int_{B^c} h_{\mathcal{B}_\varepsilon(x),B}(y) e^{-V(y)/\varepsilon} dy}{\text{cap}_{\mathcal{B}_\varepsilon(x)}(B)}$$

where

$$\text{cap}_A(B) = \varepsilon \inf_{h \in \mathcal{H}_{A,B}} \int_{(A \cup B)^c} \|\nabla h(x)\|^2 e^{-V(x)/\varepsilon} dx$$

Potential theory

$$\text{SDE: } dz_t = -\nabla V(z_t) dt + \sqrt{2\varepsilon} dW_t$$

$$\text{Equilibrium potential : } h_{A,B}(x) = \mathbb{P}^x\{\tau_A < \tau_B\}$$

Then

$$\mathbb{E}[\tau_B^x] \simeq \frac{\int_{B^c} h_{\mathcal{B}_\varepsilon(x),B}(y) e^{-V(y)/\varepsilon} dy}{\text{cap}_{\mathcal{B}_\varepsilon(x)}(B)}$$

where

$$\text{cap}_A(B) = \varepsilon \inf_{h \in \mathcal{H}_{A,B}} \int_{(A \cup B)^c} \|\nabla h(x)\|^2 e^{-V(x)/\varepsilon} dx$$

Rough a priori bounds on h show that if x potential minimum,

$$\int_{A^c} h_{\mathcal{B}_\varepsilon(x),B}(y) e^{-V(y)/\varepsilon} dy \simeq \frac{(2\pi\varepsilon)^{d/2} e^{-V(x)/\varepsilon}}{\sqrt{\det(\nabla^2 V(x))}}$$

Estimation of capacity

Truncated energy functional: retain only modes with $k \leq d$

$$\frac{1}{L}V[u] = -\frac{1}{2}z_0^2 + u_1(z_1) + \frac{1}{2} \sum_{k=2}^d \lambda_k |z_k|^2 + \dots$$

$$u_1(z_1) = \frac{1}{2}\lambda_1 z_1^2 + \frac{3}{8}z_1^4$$

Estimation of capacity

Truncated energy functional: retain only modes with $k \leq d$

$$\frac{1}{L}V[u] = -\frac{1}{2}z_0^2 + u_1(z_1) + \frac{1}{2} \sum_{k=2}^d \lambda_k |z_k|^2 + \dots$$

$$u_1(z_1) = \frac{1}{2}\lambda_1 z_1^2 + \frac{3}{8}z_1^4$$

Theorem: For all $L < \pi$,

$$\text{cap}_{\mathcal{B}_\varepsilon(u_-)}(\mathcal{B}_\varepsilon(u_+)) = \varepsilon \frac{\int_{-\infty}^{\infty} e^{-u_1(z_1)/\varepsilon} dz_1}{\sqrt{2\pi\varepsilon}} \prod_{j=2}^d \sqrt{\frac{2\pi\varepsilon}{\lambda_j}} [1 + R(\varepsilon)]$$

where $R(\varepsilon) = \mathcal{O}(\varepsilon^{1/4} |\log \varepsilon|^{5/4})$ is *uniform in d* .

Estimation of capacity

Truncated energy functional: retain only modes with $k \leq d$

$$\frac{1}{L}V[u] = -\frac{1}{2}z_0^2 + u_1(z_1) + \frac{1}{2} \sum_{k=2}^d \lambda_k |z_k|^2 + \dots$$

$$u_1(z_1) = \frac{1}{2}\lambda_1 z_1^2 + \frac{3}{8}z_1^4$$

Theorem: For all $L < \pi$,

$$\text{cap}_{\mathcal{B}_\varepsilon(u_-)}(\mathcal{B}_\varepsilon(u_+)) = \varepsilon \frac{\int_{-\infty}^{\infty} e^{-u_1(z_1)/\varepsilon} dz_1}{\sqrt{2\pi\varepsilon}} \prod_{j=2}^d \sqrt{\frac{2\pi\varepsilon}{\lambda_j}} [1 + R(\varepsilon)]$$

where $R(\varepsilon) = \mathcal{O}(\varepsilon^{1/4} |\log \varepsilon|^{5/4})$ is *uniform in d* .

Corollary:

$$\mathbb{E}_{\mathcal{B}_\varepsilon(u_-)}(\tau_{\mathcal{B}_\varepsilon(u_+)}) = \Gamma_0^{-1}(d, \varepsilon) e^{\Delta W(d)/\varepsilon} [1 + R(\varepsilon)]$$

Proposition:

Let $B = \mathcal{B}_\varepsilon(u_+)$. Assume $\sup_{d \in \mathbb{N}} \mathbb{E}[(\tau_B^d)^2], \mathbb{E}[\tau_B^2] < \infty$

Then $\mathbb{E}[\tau_B] = \Gamma_0^{-1}(\infty, \varepsilon) e^{\Delta W(\infty)/\varepsilon} [1 + \mathcal{O}(R(\varepsilon))]$

Proposition:

Let $B = \mathcal{B}_\varepsilon(u_+)$. Assume $\sup_{d \in \mathbb{N}} \mathbb{E}[(\tau_B^d)^2], \mathbb{E}[\tau_B^2] < \infty$

Then $\mathbb{E}[\tau_B] = \Gamma_0^{-1}(\infty, \varepsilon) e^{\Delta W(\infty)/\varepsilon} [1 + \mathcal{O}(R(\varepsilon))]$

Proof: Set $T_{Kr} = \Gamma_0^{-1}(\infty, \varepsilon) e^{\Delta W(\infty)/\varepsilon}$

Fix sets $B' \subset B \subset B''$ with boundaries at distance δ

Let $\Omega_{K,d} = \left\{ \sup_{t \in [0, KT_{Kr}]} \|u_t - u_t^d\|_{L^\infty} \leq \delta, \tau_{B'}^d \leq KT_{Kr} \right\}$

$$\mathbb{P}(\Omega_{K,d}^c) \leq \mathbb{P}\{Z > \delta d^\gamma\} + \frac{\mathbb{E}[\tau_{B'}^d]}{KT_{Kr}} \quad \Rightarrow \quad \lim_{d \rightarrow \infty} \mathbb{P}(\Omega_{K,d}^c) \leq \frac{2}{K}$$

Proposition:

Let $B = \mathcal{B}_\varepsilon(u_+)$. Assume $\sup_{d \in \mathbb{N}} \mathbb{E}[(\tau_B^d)^2], \mathbb{E}[\tau_B^2] < \infty$

Then $\mathbb{E}[\tau_B] = \Gamma_0^{-1}(\infty, \varepsilon) e^{\Delta W(\infty)/\varepsilon} [1 + \mathcal{O}(R(\varepsilon))]$

Proof: Set $T_{Kr} = \Gamma_0^{-1}(\infty, \varepsilon) e^{\Delta W(\infty)/\varepsilon}$

Fix sets $B' \subset B \subset B''$ with boundaries at distance δ

Let $\Omega_{K,d} = \left\{ \sup_{t \in [0, KT_{Kr}]} \|u_t - u_t^d\|_{L^\infty} \leq \delta, \tau_{B'}^d \leq KT_{Kr} \right\}$

$$\mathbb{P}(\Omega_{K,d}^c) \leq \mathbb{P}\{Z > \delta d^\gamma\} + \frac{\mathbb{E}[\tau_{B'}^d]}{KT_{Kr}} \quad \Rightarrow \quad \lim_{d \rightarrow \infty} \mathbb{P}(\Omega_{K,d}^c) \leq \frac{2}{K}$$

On $\Omega_{K,d}$ one has $\tau_{B''}^d \leq \tau_B \leq \tau_{B'}^d$

$$\Rightarrow \quad \mathbb{E}[\tau_{B''}^d] - \mathbb{E}[\tau_{B''}^d \mathbf{1}_{\Omega_{K,d}^c}] \leq \mathbb{E}[\tau_B] \leq \mathbb{E}[\tau_{B'}^d] + \mathbb{E}[\tau_B \mathbf{1}_{\Omega_{K,d}^c}]$$

Use $\mathbb{E}[\tau_B \mathbf{1}_{\Omega_{K,d}^c}] \leq \sqrt{\mathbb{E}[\tau_B^2]} \sqrt{\mathbb{P}(\Omega_{K,d}^c)}$, take $d \rightarrow \infty$ and K large enough

References

- W. G. Faris and G. Jona-Lasinio, *Large fluctuations for a nonlinear heat equation with noise*, J. Phys. A **15**, 3025–3055 (1982)
- R. S. Maier and D. L. Stein, *Droplet nucleation and domain wall motion in a bounded interval*, Phys. Rev. Lett. **87**, 270601-1 (2001)
- Anton Bovier, Michael Eckhoff, Véronique Gayrard and Markus Klein, *Metastability in reversible diffusion processes I: Sharp asymptotics for capacities and exit times*, J. Eur. Math. Soc. **6**, 399–424 (2004)
- N. B. and Barbara Gentz, *Anomalous behavior of the Kramers rate at bifurcations in classical field theories*, J. Phys. A: Math. Theor. **42**, 052001 (2009)
- N. B. and Barbara Gentz, *The Eyring–Kramers law for potentials with non-quadratic saddles*, Markov Processes Relat. Fields **16**, 549-598 (2010)
- Florent Barret, N. B. and Barbara Gentz, in preparation

Potential theory

Consider first Brownian motion $W_t^x = x + W_t$

Let $\tau_A^x = \inf\{t > 0 : W_t^x \in A\}$, $A \subset \mathbb{R}^d$

Potential theory

Consider first Brownian motion $W_t^x = x + W_t$

Let $\tau_A^x = \inf\{t > 0 : W_t^x \in A\}$, $A \subset \mathbb{R}^d$

Fact 1: $w_A(x) = \mathbb{E}[\tau_A^x]$ satisfies

$$\Delta w_A(x) = 1 \quad x \in A^c$$

$$w_A(x) = 0 \quad x \in A$$

$$G_{A^c}(x, y) \text{ Green's function} \Rightarrow w_A(x) = \int_{A^c} G_{A^c}(x, y) \, dy$$

Potential theory

Consider first Brownian motion $W_t^x = x + W_t$


Let $\tau_A^x = \inf\{t > 0 : W_t^x \in A\}$, $A \subset \mathbb{R}^d$

Fact 1: $w_A(x) = \mathbb{E}[\tau_A^x]$ satisfies

$$\Delta w_A(x) = 1 \quad x \in A^c$$

$$w_A(x) = 0 \quad x \in A$$

$G_{A^c}(x, y)$ Green's function $\Rightarrow w_A(x) = \int_{A^c} G_{A^c}(x, y) dy$



Potential theory

Fact 2: $h_{A,B}(x) = \mathbb{P}[\tau_A^x < \tau_B^x]$ satisfies

$$\Delta h_{A,B}(x) = 0 \quad x \in (A \cup B)^c$$

$$h_{A,B}(x) = 1 \quad x \in A$$

$$h_{A,B}(x) = 0 \quad x \in B$$

$$\Rightarrow h_{A,B}(x) = \int_{\partial A} G_{B^c}(x, y) \rho_{A,B}(dy)$$

Potential theory

Fact 2: $h_{A,B}(x) = \mathbb{P}[\tau_A^x < \tau_B^x]$ satisfies


$$\Delta h_{A,B}(x) = 0 \quad x \in (A \cup B)^c$$

$$h_{A,B}(x) = 1 \quad x \in A$$

$$h_{A,B}(x) = 0 \quad x \in B$$

$$\Rightarrow h_{A,B}(x) = \int_{\partial A} G_{B^c}(x, y) \rho_{A,B}(dy)$$

$\rho_{A,B}$: “surface charge density” on ∂A



Potential theory

Capacity: $\text{cap}_A(B) = \int_{\partial A} \rho_{A,B}(dy)$

Potential theory

Capacity: $\text{cap}_A(B) = \int_{\partial A} \rho_{A,B}(dy)$

Key observation: let $C = \mathcal{B}_\varepsilon(x)$, then (using $G(y, z) = G(z, y)$)

$$\begin{aligned} \int_{A^c} h_{C,A}(y) \, dy &= \int_{A^c} \int_{\partial C} G_{A^c}(y, z) \rho_{C,A}(dz) \, dy \\ &= \int_{\partial C} w_A(z) \rho_{C,A}(dz) \simeq w_A(x) \text{cap}_C(A) \end{aligned}$$

Potential theory

$$\text{Capacity: } \text{cap}_A(B) = \int_{\partial A} \rho_{A,B}(dy)$$

Key observation: let $C = \mathcal{B}_\varepsilon(x)$, then (using $G(y, z) = G(z, y)$)

$$\begin{aligned} \int_{A^c} h_{C,A}(y) \, dy &= \int_{A^c} \int_{\partial C} G_{A^c}(y, z) \rho_{C,A}(dz) \, dy \\ &= \int_{\partial C} w_A(z) \rho_{C,A}(dz) \simeq w_A(x) \text{cap}_C(A) \end{aligned}$$

$$\Rightarrow \mathbb{E}[\tau_A^x] = w_A(x) \simeq \frac{\int_{A^c} h_{\mathcal{B}_\varepsilon(x),A}(y) \, dy}{\text{cap}_{\mathcal{B}_\varepsilon(x)}(A)}$$

Potential theory

$$\text{Capacity: } \text{cap}_A(B) = \int_{\partial A} \rho_{A,B}(dy)$$

Key observation: let $C = \mathcal{B}_\varepsilon(x)$, then (using $G(y, z) = G(z, y)$)

$$\begin{aligned} \int_{A^c} h_{C,A}(y) \, dy &= \int_{A^c} \int_{\partial C} G_{A^c}(y, z) \rho_{C,A}(dz) \, dy \\ &= \int_{\partial C} w_A(z) \rho_{C,A}(dz) \simeq w_A(x) \text{cap}_C(A) \end{aligned}$$

$$\Rightarrow \quad \mathbb{E}[\tau_A^x] = w_A(x) \simeq \frac{\int_{A^c} h_{\mathcal{B}_\varepsilon(x),A}(y) \, dy}{\text{cap}_{\mathcal{B}_\varepsilon(x)}(A)}$$

Variational representation: Dirichlet form

$$\text{cap}_A(B) = \int_{(A \cup B)^c} \|\nabla h_{A,B}(x)\|^2 \, dx = \inf_{h \in \mathcal{H}_{A,B}} \int_{(A \cup B)^c} \|\nabla h(x)\|^2 \, dx$$

($\mathcal{H}_{A,B}$: set of sufficiently smooth functions satisfying b.c.)

Sketch of proof

Upper bound:

$$\text{cap} = \inf_h \Phi(h) \leq \Phi(h_+) \quad \Phi(h) = \varepsilon \int \|\nabla h(z)\|^2 e^{-V(z)/\varepsilon} dz$$

Let $\delta = \sqrt{c\varepsilon|\log \varepsilon|}$, choose

$$h_+(z) = \begin{cases} 1 & \text{for } z_0 < -\delta \\ f(z_0) & \text{for } -\delta < z_0 < \delta \\ 0 & \text{for } z_0 > \delta \end{cases}$$

where $\varepsilon f''(z_0) + \partial_{z_0} V(z_0, 0) f'(z_0) = 0$ with b.c. $f(\pm\delta) = 0, 1$

Sketch of proof

Upper bound:

$$\text{cap} = \inf_h \Phi(h) \leq \Phi(h_+) \quad \Phi(h) = \varepsilon \int \|\nabla h(z)\|^2 e^{-V(z)/\varepsilon} dz$$

Let $\delta = \sqrt{c\varepsilon|\log \varepsilon|}$, choose

$$h_+(z) = \begin{cases} 1 & \text{for } z_0 < -\delta \\ f(z_0) & \text{for } -\delta < z_0 < \delta \\ 0 & \text{for } z_0 > \delta \end{cases}$$

where $\varepsilon f''(z_0) + \partial_{z_0} V(z_0, 0) f'(z_0) = 0$ with b.c. $f(\pm\delta) = 0, 1$

Lower bound:

Bound Dirichlet Φ form below by restricting domain, taking only 1st component of gradient and use for b.c. a priori bound on h